


TILE BRITE



Mild Acid pH Disinfectant Cleaner

Product Properties

COLOR:	Translucent Blue
ODOR:	Wintergreen
VISCOSITY:	10 cps
pH:	0.7
FLASH POINT:	Not known
SPECIFIC GRAVITY:	1.050
DETERGENCY:	Excellent
SOLUBILITY IN WATER:	Complete
ACTIVE AGENT:	Quaternary ammonium chlorides

EPA REGISTERED

NOTE: Not all disinfectant and sanitizer products are available in all states.

Note: Some older tubs or sinks may be etched or discolored by this product. Test on a small hidden area first. Do not use on decorative brass fixtures, polished aluminum, natural or synthetic marble. Do not place container on Formica®.

Description

This product is a mild acid pH disinfectant cleaner which will remove hard water deposits and limestone buildups when used as directed. It quickly removes soap scum and cuts through tough grease and grime. This product inhibits the growth of mold and mildew and their odors when used as directed and improves labor results by effectively controlling odors. Use this product in hospitals, medical and dental offices, nursing homes, day care centers, nurseries, restaurants, bars, convenience stores, institutional facilities, public rest rooms, schools, colleges, athletic facilities, locker rooms, dressing rooms, shower and bath areas, exercise facilities, transportation terminals, hotels and motels, factories and business and office buildings.

Features...Benefits

One-Step Product	Improves Labor Results
Removes Hard Water Deposits.....	Improves Labor Results
Highly Effective.....	Kills Germs
Controls Mold	Controls Odors

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not for use on medical device surfaces.

PREPARATION OF DISINFECTANT USE-SOLUTION: Add 3 oz. per gallon of water (for 250 ppm water hardness), add 4 oz. per gallon of water (for 400 ppm water hardness).

DISINFECTION DIRECTIONS: Apply use-solution with a cloth, mop, sponge or sprayer or by soaking until surfaces are thoroughly wet. Treated surfaces must remain wet for 10 minutes. Wipe dry with a clean cloth or sponge or let air dry. For sprayer applications, use a coarse spray device. Spray with quick full strokes 6 - 8 inches from surface. Rub with a brush, sponge or cloth. Do not breathe spray. After use, close nozzle. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution daily or more often if use solution becomes diluted or soiled. Rinse all surfaces that come in contact with food such as countertops, appliances, tables and stovetops with potable water before reuse. This product must not be used to clean utensils, glassware and dishes.

***VIRUCIDAL DIRECTIONS:** Follow Disinfection Directions. Treated surfaces must remain wet for 1 minute. Allow surfaces to air dry.

MOLD AND MILDEW CONTROL DIRECTIONS: Add 3-4 oz. per gallon of water, to inhibit the growth of mold and mildew and their odors on hard, nonporous surfaces. Thoroughly wet all treated surfaces completely. Let air dry. Repeat application weekly or when growth or odor reappears.

TOILET BOWL/URINAL DISINFECTANT AND CLEANER DIRECTIONS: Remove gross filth before disinfection.

From concentrate: Add (3-4 oz.) to the toilet bowl or urinal and mix. Thoroughly brush or swab over exposed surfaces and under the rim with a cloth, mop, brush or sponge. Allow to stand for 10 minutes and flush.

From use-solution: Empty toilet bowl or urinal and apply a 3-4 oz. per gallon use solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer, allow to stand for 10 minutes and flush.

TO CLEAN AND DEODORIZE: Use from 1:10 dilution up to full strength. Apply product to wet the surface(s) thoroughly. Let stand for at least one minute, rinse with water or allow to air dry.

***KILLS HIV-1 ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/ OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: Clean-up must always be done wearing protective gloves, gowns, masks and eye protection.

CLEANING PROCEDURE: Blood and other body fluids containing HIV must be thoroughly cleaned from surfaces and objects before application of this product.

CONTACT TIME: Leave surface wet for 1 minute. Use a 10-minute contact time for disinfection against all other bacteria claimed.

DISPOSAL OF INFECTIOUS MATERIALS: Blood, body fluids, cleaning materials and clothing must be autoclaved and disposed of according to local regulations for infectious waste disposal.

KEEP OUT OF REACH OF CHILDREN

DANGER. Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, safety glasses or face shield), protective clothing and protective gloves (rubber or chemical resistant) when handling. Do not breathe spray. Harmful or fatal if swallowed. Avoid contamination of food. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

First Aid

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

1100 Central Industrial Dr.
St. Louis, Missouri, 63110

toll free: 800 325 9218
main: 314 771 6600
fax: 314 771 6608

email: info@interconchemical.com

www.interconchemical.com



111212-0237
© INTERCON 2011

REFER TO PRODUCT LABEL AND MSDS
FOR ADDITIONAL INFORMATION